


ALLERGY ACTION PLAN

USE 1 FORM PER CHILD FOR EACH ALLERGEN



Student _____ School _____
 DOB _____ Teacher/Grade _____
 Allergy to _____
 Asthmatic? Yes* No *Higher risk for severe reaction

STEP 1 - TREATMENT

SEND STUDENT TO HEALTH OFFICE ACCOMPANIED BY RESPONSIBLE PERSON.

The severity of symptoms can quickly change. †Potentially life threatening.

Symptoms

Give checked Medication**

***To be determined by physician authorizing treatment*

- ◆ If a student has been exposed to/ingested an allergen but has NO symptoms:
- ◆ Mouth Itching, tingling, or swelling of lips, tongue, mouth:
- ◆ Skin Hives, itchy rash, swelling of the face or extremities:
- ◆ Gut Nausea, abdominal cramps, vomiting, diarrhea:
- ◆ Throat† Tightening of throat, hoarseness, hacking cough:
- ◆ Lung† Shortness of breath, repetitive coughing, wheezing:
- ◆ Heart† Thready pulse, low blood pressure, fainting, pale, blueness:
- ◆ Other† _____ :
- ◆ If reaction is progressing, (several of the above areas affected), give:

- | | |
|--------------------------------------|--|
| <input type="checkbox"/> Epinephrine | <input type="checkbox"/> Antihistamine |
| <input type="checkbox"/> Epinephrine | <input type="checkbox"/> Antihistamine |
| <input type="checkbox"/> Epinephrine | <input type="checkbox"/> Antihistamine |
| <input type="checkbox"/> Epinephrine | <input type="checkbox"/> Antihistamine |
| <input type="checkbox"/> Epinephrine | <input type="checkbox"/> Antihistamine |
| <input type="checkbox"/> Epinephrine | <input type="checkbox"/> Antihistamine |
| <input type="checkbox"/> Epinephrine | <input type="checkbox"/> Antihistamine |
| <input type="checkbox"/> Epinephrine | <input type="checkbox"/> Antihistamine |

MEDICATION: START DATE _____ END DATE _____

Epinephrine: Inject intramuscularly.

Important; Asthma inhalers and/or antihistamines cannot be depended upon to replace epinephrine in anaphylaxis.

- Epinephrine Autoinjector 0.3mg
- Epinephrine Autoinjector 0.15mg

Antihistamine: Give _____
antihistamine/dose/route

Other: Give _____
medication/dose/route

Parent/Guardian Signature _____ Date _____

Prescriber Name _____ Phone _____

Prescriber Signature _____ Date _____

STEP 2 - EMERGENCY CALLS

PARAMEDICS (911) MUST BE CALLED IF EPIPEN OR AUVI-Q IS GIVEN. EPIPEN OR AUVI-Q ONLY LAST 15-20 MINUTES.

Call 911. State that an anaphylactic reaction has been treated, type of treatment given (i.e., EpiPen or Auvi-Q) and that additional epinephrine may be needed. Always send empty autoinjector to ER with student. Contact Parent/Guardian.

EVEN IF PARENT/GUARDIAN IS UNAVAILABLE, DO NOT HESITATE TO MEDICATE CHILD & CALL 911

EMERGENCY CONTACTS

Name	Relationship	Telephone number
1. _____	_____	_____
2. _____	_____	_____

**** Form on Page 2 to be completed ONLY if student will be carrying an Epinephrine Autoinjector ****

***** (To be completed ONLY if student will be carrying an Epinephrine Autoinjector) *****
AUTHORIZATION FOR STUDENT POSSESSION AND USE OF AN EPINEPHRINE AUTOINJECTOR
(In accordance with ORC 3313.718/8313.141)

Student name
Student address

This section must be completed and signed by the student's parent or guardian.

As the Parent/Guardian of this student, I authorize my child to possess and use an epinephrine autoinjector, as prescribed, at the school and any activity, event, or program sponsored by or in which the student's school is a participant. I understand that a school employee will immediately request assistance from an emergency medical service provider if this medication is administered. I will provide a backup dose of the medication to the school principal or nurse as required by law.

Parent/Guardian signature	Date
Parent/Guardian name	Parent/Guardian emergency telephone number ()

This section must be completed and signed by the medication prescriber.

Name and dosage of medication	
Date medication administration begins	Date medication administration ends (if known)
Circumstances for use of the epinephrine autoinjector	
Procedures for school employees if the student is unable to administer the medication or if it does not produce the expected relief	

Possible severe adverse reactions:

To the student for which it is prescribed (that should be reported to the prescriber)
To a student for which it is not prescribed who receives a dose
Special instructions

As the prescriber, I have determined that this student is capable of possessing and using this autoinjector appropriately and have provided the student with training in the proper use of the autoinjector.

Prescriber signature	Date
Prescriber name	Prescriber emergency telephone number ()

Developed in collaboration with the Ohio Association of School Nurses.
 HEA 4222 3/07